



Read the label affixed to the container for this product before applying. Use of this product according to the reference sheet is subject to the use precautions and limitations imposed by the label affixed to the container for this product. For use in hospitals and nursing homes as a disinfectant. Use in schools, colleges, commercial and industrial institutions, office buildings, hotels, motels and airports. Disinfects, cleans and deodorizes the following hard, nonporous, inanimate surfaces: floors, walls, glazed porcelain. Effective against the following pathogens:

BACTERIA	Acinetobacter baumannii ³	Listeria monocytogenes	Shigella dysenteriae	Staphylococcus aureus – Community
	Bordetella Bronchiseptica	Lactobacillus casei	Shigella flexneri Type 2	Associated Methicillin-Resistant - CA-MRSA (USA400)
	Cornybacterium ammoniagenes	Pasteurella multocida	Shigella sonnei	Staphylococcus aureus – VISA
	Enterobacter aerogenes	Proteus mirabilis	Staphylococcus aureus	Staphylococcus epidermidis ¹
	Enterobacter cloacae	Proteus vulgaris	Staphylococcus aureus ¹	Streptococcus pyogenes
	Enterococcus faecalis – (VRE)	Pseudomonas fluorescens	Staphylococcus aureus - MRSA	Streptococcus pyogenes group A
	Enterococcus faecalis	Pseudomonas tolaasii	Staphylococcus epidermidis	Xanthomonas maltophilia ¹
	Enterococcus faecalis ¹	Pseudomonas aeruginosa ¹	Staphylococcus aureus – Community	
	Escherichia coli ¹	Salmonella enterica	Associated Methicillin-Resistant - CA-MRSA (USA300)	
	Fusobacterium necrophorum	Salmonella typhi		
	Klebsiella pneumoniae ¹	Serratia marcescens		
	¹ Clinical Isolate	² Clinical Isolate (Fairfax Hospital) Resistant – Ampicillin, Bactrim, Cefazolin, Cefazdime, Ceftriaxone, Ciprofloxacin, Gentamicin, Levofloxacin, Tobramycin	³ subsp. – serotype paratyphi B, pullorum, typhi, thyphimurium, enteritidis	

VIRUSES	*Hepatitis B virus (HBV)	*Influenza A virus / Hong Kong
	*Hepatitis C virus (HCV)	*Pandemic 2009 H1N1 Influenza A virus
	*Herpes Simplex virus Type 1	*Paramyxovirus
	*Herpes Simplex virus Type 2	*Rotavirus
	*Human coronavirus	*SARS Associated Coronavirus
	*HIV-1 (AIDS virus)	*Vaccinia virus
	*HIV-2	

ANIMAL VIRUSES	*Avian Influenza A virus	*Infectious bovine rhinotracheitis virus
	*Avian Influenza (H5N1)	*Infectious bronchitis virus (Avian IBV)
	*Canine distemper virus	*Pseudorabies virus
	*Feline leukemia virus	*Rabies virus
	*Feline picornavirus	

OTHER	Fungi: Candida albicans	Trichophyton mentagrophytes
	Mildew: Aspergillus niger	

ACTIVE INGREDIENTS	Octyl decyl dimethyl ammonium chloride.....	6.510%
	Diocetyl dimethyl ammonium chloride	3.255%
	Didecyl dimethyl ammonium chloride.....	3.255%
	n-Alkyl C ₁₄ 50%, C ₁₂ 40% and C ₁₆ 10%) dimethyl benzyl ammonium chloride.....	8.680%
	INERT INGREDIENTS:	78.300%
	TOTAL:	100.000%

DILUTION	1:256
	½ oz. per gallon of water

KEEP OUT OF REACH OF CHILDREN DANGER

ADDITIONAL DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Non-Acid Toilet Bowl Disinfection/Cleaner Directions: Remove gross filth prior to disinfection.

From concentrate: Add ½ ounce per gallon of water to the toilet bowl and mix. Brush thoroughly over exposed surfaces and under the rim, allow to stand for 10 minutes and flush.

From use-solution: Empty toilet bowl or urinal and apply ½ ounce per gallon of water use-solution to exposed surfaces, brush or swab thoroughly under the rim with a cloth, mop, sponge or sprayer, allow to stand for 10 minutes and flush.

For heavy duty use: Empty toilet bowl or urinal and apply 2 ounces per gallon of water use-solution to exposed surfaces, brush or swab thoroughly under the rim with a cloth, mop, sponge or sprayer, allow to stand for 10 minutes and flush.

VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS DISINFECTION DIRECTIONS: For cleaning and disinfecting hard nonporous surfaces: equipment used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of 1/2 oz. per gallon of water (or equivalent dilution) for a period of 10 minutes. Immerse all animal handling and restraining equipment as well as forks, shovels and scrapers used to remove litter and manure. Wipe or allow to air dry. Ventilate buildings and other closed spaces. Thoroughly scrub all treated surfaces that come into contact with food, including equipment used for feeding or watering, with soap or detergent and rinse with potable water before reuse. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

Poultry (And Swine) Premise: Remove all animals and feeds from premises, trucks, cars, coops and crates. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Use ½ ounce of this product per gallon of water. Saturate surfaces with the disinfecting solution for a period of 10 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.

To Control Mold and Mildew on Large, Inflatable, Non-Porous Plastic and Rubber Structures (animals, promotional items, moonwalk, slides, obstacle course play equipment, exercise equipment, and wrestling mats): Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of ½ ounce of this product per 1 gallon of water for a period of 10 minutes. Ventilate buildings and other close spaces. Do not use equipment until treatment has been absorbed, set or dried.

SKUs- 2/3100 mL (5116791) and 55 gal. drum (5501584)